
Friday 05.02.2021


Published 04 02 2021, 17:00



AM




PM







Danger Level 3 - Considerable



As the day progresses, wet and gliding avalanches are possible. Weakly bonded old snow requires caution.

The avalanche danger will increase a little during the day. As a consequence of warming during the day and solar radiation more frequent wet loose snow avalanches are to be expected. This applies on extremely steep sunny slopes. Avalanches can also penetrate deep layers and reach dangerously large size. In addition a substantial danger of gliding avalanches exists. Areas with glide cracks are to be avoided. Weak layers in the old snowpack can still be released by individual winter sport participants. Caution is to be exercised in all aspects above approximately 1900 m, especially on very steep slopes, as well as at transitions from a shallow to a deep snowpack.

In addition the fresh wind slabs in high Alpine regions are prone to triggering in some cases, especially adjacent to ridgelines on shady slopes.

Experience in the assessment of avalanche danger is required.

Snowpack

Danger patterns dp.10: springtime scenario dp.7: snow-poor zones in snow-rich surrounding

The spring-like weather conditions will give rise to increasing moistening of the snowpack. Outgoing longwave radiation during the night will be reduced. The surface of the snowpack is frozen, but not to a significant depth and will soften during the day. This applies in all aspects at low and intermediate altitudes, as well as on very steep sunny slopes also at elevated altitudes.

Faceted weak layers exist in the centre of the snowpack. This applies in particular above approximately

Towards its base, the snowpack is largely stable.

The fresh wind slabs are lying on soft layers in high Alpine regions. This applies on shady slopes.

Avalanche.report

Friday 05.02.2021

Published 04 02 2021, 17:00




Tendency

The avalanche danger will increase during the day.



Danger Level 3 - Considerable



As a consequence of warming during the day and solar radiation more frequent wet and gliding avalanches are possible. In some places avalanches can be released in the weakly bonded old snow and reach large size.

The avalanche danger will increase a little during the day. As a consequence of warming during the day and solar radiation more frequent wet loose snow avalanches are to be expected. This applies on extremely steep sunny slopes. Avalanches can also penetrate deep layers and reach dangerously large size. An appreciable danger of gliding avalanches exists. This applies in particular in the west and in the northwest. Areas with glide cracks are to be avoided.

Dry avalanches can additionally be released in the weakly bonded old snow by a single winter sport participant. This applies above approximately 1900 m, especially in areas where the snow cover is rather shallow, as well as at transitions from a shallow to a deep snowpack. Between approximately 1900 and 2300 m the avalanche prone locations are more prevalent and the danger is slightly greater. Avalanches can penetrate deep layers and reach dangerously large size. Remotely triggered avalanches are possible. Experience and restraint are required.

In particular in the vicinity of peaks sometimes avalanche prone wind slabs formed.

Snowpack

Danger patterns

(dp.7: snow-poor zones in snow-rich surrounding)

dp.10: springtime scenario

The spring-like weather conditions will give rise to increasing moistening of the snowpack, especially at low and intermediate altitudes, as well as on very steep sunny slopes.

Avalanche prone weak layers exist in the centre of the snowpack, especially between approximately 1900 and 2300 m in all aspects. Released avalanches and stability tests confirm the existence of a weak snowack.

Avalanche.report

Friday 05.02.2021

Published 04 02 2021, 17:00



Tendency

The danger of wet and gliding avalanches will increase during the day.



Danger Level 2 - Moderate





Tendency: Constant avalanche danger on Saturday 06 02 2021





weak layer







As a consequence of warming during the day and solar radiation mostly small moist snow slides are possible as the day progresses.

As a consequence of warming during the day and solar radiation individual moist avalanches are possible as the day progresses, but they will be mostly small. Caution is to be exercised in particular on extremely steep sunny slopes.

Weak layers in the old snowpack can still be released in some places by individual winter sport participants in particular above approximately 1900 m.

Snowpack

Danger patterns

dp.7: snow-poor zones in snow-rich surrounding

dp.10: springtime scenario

Avalanche prone weak layers exist in the centre of the snowpack. This applies in particular above approximately 1900 m. The spring-like weather conditions will give rise to gradual moistening of the snowpack. The snowpack will be moist at intermediate altitudes. The snowpack will be wet all the way through at low altitude.

Tendency

Increase in danger of moist and wet avalanches.