

# Weakly bonded old snow is to be evaluated with care and prudence. Increase in avalanche danger as a consequence of solar radiation.

Weak layers in the old snowpack can still be released in very isolated cases, this applies even in case of a single winter sport participant. Caution is to be exercised in particular on very steep shady slopes above approximately 2200 m, as well as on very steep east facing slopes above approximately 2400 m, especially on little used northeast and east facing slopes.

Night: As a consequence of the rain small natural wet avalanches are possible below approximately 2200 m. This applies in particular on shady slopes.

As the day progresses the likelihood of wet avalanches being released will increase. As a consequence of solar radiation, the natural avalanche activity will increase. This applies in particular on very steep east, south and west facing slopes below approximately 2400 m.

#### Snowpack

Danger patterns

dp.1: deep persistent weak layer

Up to 2000 m rain will fall in the next few hours. In some localities up to 15 cm of snow will fall above approximately 2000 m.

Faceted weak layers exist in the old snowpack, especially on shady slopes above approximately 2200 m, as well as on east and west facing slopes above approximately 2400 m.


Outgoing longwave radiation during the night will be reduced in some case. The surface of the snowpack is frozen, but not to a significant depth and will soften during the day, in particular at intermediate altitudes. Hardly any snow is lying at low altitude. The snowpack will be moist at low and intermediate altitudes.

# Tendency

Tuesday: Hardly any increase in avalanche danger as a consequence of warming during the day.







### Weakly bonded old snow is to be evaluated with care and prudence.

Weak layers in the old snowpack can still be released in very isolated cases, this applies even in case of a single winter sport participant. Caution is to be exercised in particular on very steep shady slopes above approximately 2200 m, as well as on very steep east facing slopes above approximately 2400 m, especially on little used northeast and east facing slopes.

Night: As a consequence of the rain small natural wet avalanches are possible below approximately 2200 m. This applies in particular on shady slopes.

As the day progresses the likelihood of wet avalanches being released will increase a little. This applies in particular on very steep east, south and west facing slopes below approximately 2400 m.

#### Snowpack

Danger patterns

dp.1: deep persistent weak layer

Up to 2000 m rain will fall in the next few hours. In some localities up to 15 cm of snow will fall above approximately 2000 m.

Faceted weak layers exist in the old snowpack, especially on shady slopes above approximately 2200 m, as well as on east and west facing slopes above approximately 2400 m.


Outgoing longwave radiation during the night will be reduced in some case. The surface of the snowpack is frozen, but not to a significant depth and will soften during the day, in particular at intermediate altitudes. Hardly any snow is lying at low altitude.

### Tendency

Tuesday: Hardly any increase in avalanche danger as a consequence of warming during the day.







## Weakly bonded old snow is to be evaluated with care and prudence.

Weak layers in the old snowpack can still be released by individual winter sport participants. This applies in particular on very steep shady slopes above approximately 2200 m, as well as on very steep east and west facing slopes above approximately 2400 m. Caution is to be exercised on little used shady slopes. In addition the older wind slabs at high altitudes and in high Alpine regions are capable of being triggered

in isolated cases still.

As the day progresses the likelihood of wet avalanches being released will increase, in particular on sunny slopes below approximately 2600 m. Backcountry tours should be concluded timely.

#### Snowpack

Danger patterns

dp.1: deep persistent weak layer

Faceted weak layers exist in the old snowpack, especially on shady slopes above approximately 2200 m, as well as on east and west facing slopes above approximately 2400 m.

The old wind slabs are in individual cases still prone to triggering at high altitudes and in high Alpine regions.

Outgoing longwave radiation during the night will be reduced in some case. The surface of the snowpack is frozen, but not to a significant depth and will soften during the day. The weather conditions will bring about a slight weakening of the snowpack as the day progresses.

# Tendency

The danger of dry avalanches will decrease gradually. As the day progresses as a consequence of warming during the day and solar radiation there will be only a slight increase in the danger of wet avalanches.





### Danger Level 1 - Low



Tendency: Constant avalanche danger  $\longrightarrow$  on Tuesday 21 03 2023

## Currently there are favourable avalanche conditions mostly.

Individual avalanche prone locations are to be found in particular on steep, little used shady slopes at high altitude.

### Snowpack

The snowpack will be generally stable. Only a small amount of snow is lying for the time of year at low and intermediate altitudes. Some rain will fall in some localities. Above approximately 2000 m snow will fall.

## Tendency

Low avalanche danger will persist.

